

2025



Missouri Black Bear Program Annual Report



Missouri Department of Conservation

Prepared by:

Nate Bowersock – Resource Scientist

Contributors:

Nate Bowersock and Lakelen Murphy

Cover photo: Photo of research bear 2324 with her 2 cubs in 2025.

Table of Contents

2025 Missouri Black Bear Program Annual Report	1
Black Bear Program Mission	2
Introduction	3
Black Bear History in Missouri	3
Black Bear Management Plans	4
Purpose Statement	4
Using Science to Manage a Self-Sustaining Population	5
Previous Research	5
Current Research	5
Bear Sightings Reports	8
Hunting.....	9
Season Parameters	9
Harvest Report	9
Increasing Public Awareness	10
Outreach and Public Engagement	10
Black Bear Hunting and Ecology Classes	10
Addressing and Mitigating Stakeholder Conflicts	11
Bear Wildlife Damage Response	12
Acknowledgements and Publications.....	13
Acknowledgments	13
Publications (from 2010-present)	13

Black Bear Program Mission

The mission of Missouri Department of Conservation's (MDC) Black Bear Program is to use science-based methods to manage a self-sustaining population of black bear (a native species), increase bear awareness, minimize human-bear conflict, and provide recreational opportunities for all Missourians. To enact this mission, the Black Bear Program is guided by three management goals:


Goal 1) Bear Research and Monitoring - Use science-based methods to manage a self-sustaining population of black bear, focusing on research and monitoring, population management, and habitat management.

Goal 2) Education and Outreach - Increase statewide awareness of Missouri's black bear population and management program through coordinated outreach and public education.

Goal 3) Human Bear Conflict - Minimize and address human-bear conflicts.

The Black Bear Program is managed by MDC's Science Branch Black Bear/ Furbearer Biologist in collaboration with many MDC staff, who are responsible for monitoring and managing the state's black bear population and developing annual regulation recommendations based on: annual bear sighting reports, harvest data, hunter surveys, MDC staff surveys, public comments, population models, and ongoing research studies. The conservation of Missouri's black bear population along with all other wildlife species is made possible by the one-eighth of one percent Conservation Sales Tax, permit sales, and income generated by fish and wildlife tourism. In addition, conservation efforts would not be possible without the help of a wide range of MDC staff, university researchers, Missouri citizens, and support from local landowners. The bear program would not be able to complete the mission set forth by the MDC without this assistance.

Thank you!!!



Equal Opportunity to Participate

Equal opportunity to participate in, and benefit from, programs of the Missouri Department of Conservation is available to all individuals without regard to their race, color, nationality, sex, age, or disability. Questions should be directed to the Department of Conservation, PO Box 180, Jefferson City, MO 65102, 573-751-4115 (voice) or 800-735-2966 (TTY), or to the U.S. Fish and Wildlife Service Division of Federal Assistance, 4401 N. Fairfax Drive, Mail Stop: MBSP-4020, Arlington, VA 22203.

Introduction

Black Bear History in Missouri

Black bears were historically abundant in Missouri but became seemingly nonexistent by the early 1900s. Many early county histories contain notes and reports of the remarkable number of bears in all areas of the state. Bears were a staple item for early settlers and were widely used for food as well as for their fat and skin. In fact, bears were more commonly harvested by pioneers and early travelers than any large mammal, other than deer. However, by the 1830s and 1840s, bears were rare in northern Missouri, and by 1894, bears were almost extirpated from the Ozarks. Bears were still present in southeastern Missouri in the 1920s, however by the 1930s and '40s, few bears could be found in the state.



Black Bear illustration Courtesy of MDC.


distribution since that time, and thereafter, sightings of black bear began to rise in Missouri, likely from bears crossing into Missouri from Arkansas.

The Arkansas Game and Fish Commission initiated a black bear restoration program in 1958. From 1958 to 1968, 254 black bears were captured in Minnesota and Manitoba, Canada and released in the Ozark and Ouachita mountains of western Arkansas. The Arkansas bear population expanded in both size and

From 1950–1972, there were 54 reported occurrences of bears in 27 Missouri counties. In 1990, a request for sighting information published in the June Conservationist magazine resulted in 55 reports of sightings in 26 counties. Over the last two decades, reports have increased substantially. From 2000–2010, MDC received 512 bear reports in 75 counties. Intensive research studies including Missouri's Black Bear Hair Snare study and collaring study were started in 2010. These studies have allowed for better understanding and tracking of the bear population. In 2025 alone, MDC received 372 bear reports in 70 counties. Overall, black bears have been reported in 107 counties, although it should be noted that reports are not always verified and likely include some percent that are misidentified, but in general, the distribution and number of bear reports is steadily increasing. This demonstrates that the once nearly extirpated bear population is successfully recolonizing the state.

Black Bear Management Plans

To guide the continued conservation and management of Missouri black bear population, a black bear management plan was first developed in 1993 by MDC, in partnership with the Missouri Department of Natural Resources, the National Park Service, and the U.S. Forest Service. As the black bear population continued to grow and conservation objectives changed, new management plans were developed in 2008 and most recently in 2020. The plans identified critical components of a multifaceted approach to bear management in Missouri with the goal of providing a strategic direction for the MDC bear management program to follow. To stay on top of changes to the bear population and conservation efforts, new management plans will be generated every 10 years. The 2020-2030 Missouri Bear Management Plan can be found on the MDC website [Missouri Bear Management Plan \(2020-2030\)](#).



Trail camera photo from the Missouri Black Bear Research Project. Howell County, 2017.

Purpose Statement

This annual report summarizes the efforts completed by the Missouri Department of Conservation Black Bear Program in pursuit of managing a self-sustaining black bear population in Missouri. The scientific study and management of black bears in Missouri are overseen by the Missouri Department of Conservation Black Bear Program, housed within the Game and Regulations Unit of the Science Branch. The black bear program is headed by the State Black Bear and Furbearer Biologist and is advised by an inter-branch Black Bear Working Group. All regulatory actions must be proposed and ultimately approved by the Conservation Commission. The following sections include updates on black bear research, the 2025 harvest season, and other management activities throughout the state.

Questions regarding the content of this report can be directed to Nate Bowersock at Nathaniel.Bowersock@mdc.mo.gov or visit <https://mdc.mo.gov/discover-nature/field-guide/american-black-bear>.

Using Science to Manage a Self-Sustaining Population

Previous Research

Starting in 2010, research began to further MDC's understanding of the expanding black bear population in Missouri. To estimate the bear population at the time, a hair snare study was conducted. Hair snares are barbed-wire enclosures that are baited with a scent lure to encourage bears to crawl over or under a single strand of barbed-wire, with the goal of collecting hair from bears. The hair samples collected from snares were used to identify unique individual bears with DNA analysis and was used in population models. From this initial population study, it was estimated there were around 300 bears in Missouri in 2012.



Black bear hair that was caught on a barbed wire hair snare.

Current Research

Survival, Recruitment, and Movement

Due to black bears' cryptic nature and the dense forest cover they prefer to inhabit; focal studies of bears can be quite difficult to conduct. Therefore, researchers have found that much can be learned about bears and other wildlife by capturing them and fitting them with radio tracking collars. Starting in 2010, MDC in conjunction with university researchers, began capturing bears

using large box traps and fitting them with GPS radio tracking collars. When bears are captured, they are chemically immobilized and sedated so they can be safely handled by MDC staff. Once a bear is immobilized, a series of biological measurements (i.e., body weight and length) and samples (i.e., hair and blood) are collected. In addition, each bear is marked with both a set of uniquely numbered ear tags and a PIT tag (microchip) so bears can be identified if they are seen or captured again. Capture operations have been conducted continually for over a decade and a half now as part of the long-term research and monitoring objectives of the bear program. Data collected from GPS radio



Trail camera photo of a bear examining a research bear trap.

collars has been used to study bear survival, recruitment, and movement rates. As of 2025, 208 black bears (184 female and 24 males) have been collared and the data collected from these animals has helped MDC better understand the ecology of bears in Missouri.

Trapping – In 2025, a total of 22 individual bears (11 females, 11 males) were captured. The team handled 17 bears (8 females, 9 males) and collared 7 females. Six of the collared females

were unmarked and one was previously marked. The 2025 trapping season started early June and continued through early July. The rate of bears trapped daily was fairly consistent with 1-2 being captured on average daily, and a high for the season being 5 in one day. However, the rate of capture had a slight increase in July as the damage biologists used dynamic trapping methods in efforts to deploy all collars.

Denning – To study the recruitment rates of bears (number of young born each year compared to how many survive within a year), MDC tracked GPS collared females to their dens to count the number of cubs they have each year. When possible, staff attempted to sedate females to safely enter dens to check the physical condition of both the collared females and their cubs. Cubs found at dens were measured, had genetic samples taken, and marked with PIT tags so they could be identified in the future if found through capture or mortality events. If dens could not be safely entered, females were left alone, and trail cameras were set up outside of dens to count the number of cubs that emerge. In the following year, MDC staff try to relocate females to count the number of yearling cubs they have to assess the survival rate of cubs. The survival estimates gained from this study are used in population studies and help assess the overall health of the bear population.

Due to the premature drop of two collars deployed in summer 2024 and the timing of refurbished collars' arrival, we only had two collared sows to potentially work the 2025 denning season. Only one of the dens was worked due to accessibility, and 2 cubs (1 female and 1 male) were marked. Overall, MDC has visited 245 dens, counted 176 cubs, and marked 152 cubs.



Black bear crew collecting data on collared bear 2324 at her den in early 2024. Photos courtesy of William Harris – MDC.

Bear Mortalities – MDC responded to a total of 14 confirmed bear mortalities in 2025, including those legally harvested by hunters. Only three of the bears were previously marked research bears, the remaining were unmarked. Two of the mortalities were caused by vehicle collisions, one bear was poached, one was the result of pet conflict, and one bear was euthanized due to severe mange infestation. Nine bears were legally harvested by hunters.

Abundance and growth rate estimates - Using the initial population estimate of 300 black bears in 2012, along with data collected from collared bears such as their annual survival and reproductive rates, a population model is used to track trends in the bear population over time (Figure 1). In 2025, we estimated the black bear population to be approximately 1,065 black bears in the state with an annual growth rate of 9%, which indicates continual growth of the bear population. The increase in bear numbers is also reflected in the increased number of bear sightings we have received, including increasing sightings outside of the core black bear range in southern Missouri.




Figure 1. Annual black bear population abundance estimates for Missouri (orange) based on predictive population models. Estimates based on the initial population estimate of 300 bears in 2012 and upper and lower confidence limit bars of population trend lines.

Population abundance and density distribution - The 2-year hair-snare study that started in 2022 wrapped up data collection in the summer of 2023. Over 500 hair snares were deployed across 48 sampling sites, and more than 1,000 hair samples were collected. Hair samples were collected at all but 9 sampling sites. The goal of the study is to generate a new population abundance estimate for Missouri. In addition, this study will explore questions looking into how landscape features might influence the density distribution of bears to further our understanding of how bears are distributed across the state to help focus future management efforts. Due to some unexpected delays results from this study will likely not be available until summer 2026.

Other Missouri related research - Genetic samples taken from Missouri black bears were used to assess the color variation seen in American black bears found across North America. In Missouri, black bear coat color can range from black to blonde, with dark and reddish brown (cinnamon) variations also seen. Results from this study found that Missouri black bears traditionally had black coats, but after the population restoration work that Arkansas conducted in the 1960's, bringing

bears from Minnesota and Canada down to the region, brown and other coat color genetic variations were introduced into region which explains the broad coat color variation in Missouri.

Bear Sightings Reports

The MDC has been collecting reports of bear sightings throughout the state of Missouri since the late 1980’s (Figure 2). These reports include in-person sightings, photos captured on remotely triggered cameras, and any evidence of bear activity (e.g., tracks, scat, or damage). These reports have been pivotal to tracking the expansion of the bear population over the last several decades.




Figure 2. Count of bear sightings reported to MDC since 1987 – 2025.

Maintaining a record of bear sightings helps MDC understand trends in bear observations over time. These trends are especially useful in tracking human-bear conflicts to focus public outreach and mitigative actions. Reports are collected through an online reporting system, which can be found at mdc.mo.gov/reportbears.

These reports are then cataloged and categorized by Black Bear Program staff. In 2025, the Department received 332 sighting reports from members of the public, MDC staff, and partnering agencies throughout the state. Sightings were reported in 69 (61%) of Missouri counties (Figure 3).




Figure 3. Maps of the black bear sightings by county in 2025. The black line that crosses the state from the SW corner to the eastern portion of the central region depicts Interstate 44.

Hunting

Season Parameters

The 2025 bear hunting season began the third Saturday in October and ran for 14 consecutive days from October 18-31 within 3 Bear Management Zones (BMZ: Figure 5). A total of 600 permits (BMZ 1: 300, BMZ 2: 225, BMZ 3: 75) were made available, with 10% of permits in each zone being allocated to qualifying resident landowners whose property was within the zone they applied for.

Harvest quotas for each BMZ were determined based on black bear population growth and harvest simulations, and a 10% hunter harvest success rate, with caps on harvest being set as follows for each zone: BMZ 1 – 20 bears, BMZ 2 – 15 bears, BMZ 3 – 5 bears. These values are not a target harvest, but a cap on harvest that allows for maximum hunter opportunity while preventing impact on population growth.

All methods of harvest acceptable for deer are permitted for black bear except for atlatl. In addition, the use of dogs and baiting were not permitted, consistent with other large game hunting regulations in the state. Those awarded bear hunting permits were not allowed to be assisted by another individual during their hunt, unless the other individual(s) had also drawn a permit (with exceptions for minors and those with disabilities requiring assistance).

Harvest Report

In 2025, a total of 5,249 (BMZ 1: 2,183, BMZ 2: 2,050, BMZ 3: 1,016) Missouri residents applied for a bear permit, and of the 600 permits awarded, 487 were purchased. At the end of the season, a total of 9 bears (7 females, 2 males) were harvested (Figure 4), 7 in BMZ1 and 2 in BMZ2. Of the bears taken, all were harvested using firearms. The age range of harvest animals was 1.5 to 9.5 years and two of the bears harvested were marked during capture operations in 2022 and 2024.




Figure 4. Map displaying Missouri’s bear management zones and the number of bears harvested per county in each zone in 2025.

Following each hunting season, a post-season survey is sent out to all hunters that were drawn to assess overall effort and satisfaction. Questions include asking hunters about where they hunted, if they were able to find bear sign before or during the bear season, did hunters see bears while hunting, overall hunter satisfaction with the hunting season, and more. Of the hunters that responded to the survey, 42% of hunters attended a bear hunting class in 2025, which was similar to attendance in 2024 (41%). Unlike last year, less than half of hunters across all BMZs found bear sign before going out to hunt, and few hunters (7% of respondents) saw a bear while hunting that they did not harvest.




Figure 5. Post-season survey results regarding hunter satisfaction with the bear hunting experience from the 2025 hunting season.

Overall, satisfaction rates between BMZs in 2025 were comparable (Figure 5). Additionally, hunters who reported spending more than 20 days scouting had a significantly higher bear harvest success rate compared to those who spent fewer days scouting.

Increasing Public Awareness

Outreach and Public Engagement

We continued to engage the public to increase their awareness of our ever growing and expanding black bear population through a series of educational programs, news articles, radio interviews, and podcasts. Our hope is that as more people become aware that Missouri is bear country, citizens can learn how to live responsibly with bears.

Black Bear Hunting and Ecology Classes

Since the establishment of Missouri black bear hunting season, MDC has put on bear hunting classes (Black Bear Hunting: Basics) for those that might be interested in learning more about Missouri’s black bear season. Attendees learned about wildlife identification, habits and habitats, regulations, safety, hunting strategies, equipment, and game care. Furthermore, MDC also held a more in-depth training class (Black Bear Hunting Beyond the Basics) for those individuals that were drawn for the opportunity to purchase a bear hunting permit. This class touched on the same topics but expanded on a number of hunting related topics such as scouting, hunting logistics, and time for Q&A for hunters.

Be BearWise

As the bear population started to grow and expand across the state, MDC saw a need to begin intensive educational and outreach activities to help Missourians learn how to live with bears, which led to the development of the “Be Bear Aware” program. However, in recent years Missouri has joined the BearWise program, a nation-wide education program developed by black bear biologists and supported by state wildlife agencies, such as MDC, that provides consistent messaging about how to live responsibly with bears—whether you live in Missouri or are going on vacation to Montana or Tennessee. BearWise shares ways to prevent conflicts and encourages community initiatives to keep bears wild. You can learn more about this program at <https://bearwise.org>, or visit MDC’s website: <https://mdc.mo.gov/bearwise>.

At Home BearWise Basics

1. Never feed or approach bears
2. Secure food, garbage and recycling
3. Remove bird feeders when bears are active
4. Never leave pet food outdoors
5. Clean and store grills, smokers
6. Alert neighbors to bear activity



If you see a bear in Missouri or find bear sign, we encourage the public to report these sightings to MDC at mdc.mo.gov/reportbears. In addition, if you are regularly seeing bear activity on your property or are experience conflict with a bear, it is important to contact your MDC Regional Office, local Conservation Agent, or regional Damage Biologist.

Addressing and Mitigating Stakeholder Conflicts



A trail camera photo of a black bear attempting to access an unsecured dumpster in southern Missouri.

Black bears are an important species in Missouri for their intrinsic value and their contributions to local biological diversity. However, sometimes bears may be considered a problem by some members of the public when involved in human-bear conflicts. These interactions are likely to increase over time as the black bear population and their geographic distribution expands. For the continuation of our healthy bear population and assuring public safety, it is imperative to deploy a comprehensive strategy addressing and mitigating human-bear conflicts across their range.

Bear Wildlife Damage Response

MDC’s Wildlife Damage Management Program employs a team of Wildlife Damage Biologists, or Human-Wildlife Interaction Specialists, that provide technical information and assistance with wildlife damage prevention and control. These damage biologists use flyers, door hangers, and one-on-one interactions from advising on securing trash or waste, to assisting in putting up electric fencing and other deterrents to address conflict issues and prevention. Human bear conflicts are most common outside the bear hibernation periods and typically peak in the early summer. Since 2010, damage biologists have responded to a long list of conflict issues, most of which related to food storage or livestock (such as chicken coops or beehives) related issues (Figure 6).




Figure 6. The proportion of human-bear conflicts reported to MDC from 1991 – 2025. The other category includes events such as vehicle collisions or bears in city/town.

In 2025, a similar number of human-bear conflicts were reported when compared to previous years, with trash being the most common issue, followed by feed storage and livestock and beehives (Figure 7). To help deal with these issues, damage biologists regularly recommend removing any food attractants from areas of conflict and secure trash cans and other attractants with electric fencing. These resolutions help mitigate many conflict issues.




Figure 7. The proportion of reported bear–community interaction during 2025 in Missouri related to sources of food or direct interactions with property, pets or livestock, trash or feed of different types.

Acknowledgements and Publications

Acknowledgments

The Program recognizes the efforts by all its collaborators and would like to extend the utmost appreciation towards the stated and unstated herein. These include countless Missouri landowners that graciously allow the Program access to their property for research activities. Department staff and external agency partners (e.g., U.S. Forest Service, USFWS Wildlife Restoration Program) provide exceptional service to the Program throughout the year. University collaborators (e.g., the State University of New York, Mississippi State University, Michigan State University) collaborate with MDC on the high-quality research that informs management decisions. Thank you!

Publications (from 2010-present)

From the research conducted starting in 2010, more than 23 peer reviewed publications were published in scientific journals. We hope to continue to share our results of future studies in scientific publications as these studies are completed.

Al-Warid, HS, Beringer J, Hiller TL, Belant, JL, Gomper, ME. 2016. Community composition of Ixodid ticks parasitizing American black bears in Missouri, USA. *Ursus* 27:61-66.

Boudreau, M. R., Gantchoff, M. G., Ramirez-Reyes, C., Conlee, L., Belant, J. L., & Iglay, R. B. (2021). Using habitat suitability and landscape connectivity in the spatial prioritization of public outreach and management during carnivore recolonization. *Journal of Applied Ecology*, 59(3), 757-767.

Boudreau, M. R., Gantchoff, M. G., Conlee, L., Anderson, C., Bowersock, N. R., Belant, J. L., & Iglay, R. B. (2024). A harvest framework for a recovering American black bear population. *The Journal of Wildlife Management*, 88(1), e22508.

Duquette JF, Belant JL, Wilton CM, Fowler N, Waller BW, Beyer Jr DE, Svoboda NJ, Simek SL, Beringer J. 2017. Black bear (*Ursus americanus*) functional resource selection relative to intraspecific competition and human risk. *Canadian Journal of Zoology* 95:203–12.

Gantchoff MG, Belant JL. 2017. Regional connectivity for recolonizing American black bears (*Ursus americanus*) in southcentral USA. *Biological Conservation* 214:66–75.

Gantchoff MG, Conlee L, Belant JL. 2020. Planning for carnivore recolonization by mapping sex-specific landscape connectivity. *Global Ecology and Conservation* 21: e00869.

Gantchoff MG, Conlee L, Belant JL. 2019. Conservation implications of sex-specific landscape suitability for a large generalist carnivore. *Diversity and Distributions* 25:1488–1496.

Gantchoff MG, Beyer Jr D, Belant JL. 2019. Reproductive class influences risk tolerance during denning and spring for American black bears (*Ursus americanus*). *Ecosphere* 10: e02705.

Gantchoff MG, Hill JE, Kellner KF, Fowler NL, Petroelje TR, Conlee L, Beyer Jr DE, Belant JL. 2020. Mortality of a large wide-ranging mammal largely caused by anthropogenic activities. *Scientific Reports*: 10, 8498.

Gantchoff MG, Wang G, Beyer D, Belant JL. 2019. Scale-dependent home range optimality for a solitary omnivore. *Ecology and Evolution* 23:12271–12282.

Gantchoff, M. G., Conlee, L., Boudreau, M. R., Iglay, R. B., Anderson, C., & Belant, J. L. (2022). Spatially-explicit population modeling to predict large carnivore recovery and expansion. *Ecological Modelling*, 470, 110033.

Gantchoff, M. G., Conlee, L., and Belant, J. (2022). The effectiveness of opportunistic public reports versus professional data to estimate large carnivore distribution. *Ecosphere*, 13(2), e3938.

Hiller TL, Belant JL, Beringer J, Tyre AJ. 2015. Resource selection by recolonizing American black bears in a fragmented forest landscape. *Ursus* 26:116–128.

Hiller TL, Belant JL, Beringer J, Tyre AJ. 2015. Sexual size dimorphism mediates effects of spatial resource variability on American black bear space use. *Journal of Zoology* 296:200–207.

Hiller TL, Belant JL, Beringer J, Tyre AJ. 2017. Shape complexity of space used by American black bears influenced by sex and intensity of use. *Basic and Applied Ecology* 18:67–74.

Kristenson, T.V., Puckett, E.E., Landguth, E.L., Belant, J.L., Hast, J.T., Carpenter, C., Sajeck, J.L., Beringer, J., Means, M., Cox, J.J., Eggert, L.S., White, D., Jr., and Smith, K.G. 2018. Plasticity of dispersal in black bears (*Ursus americanus*) in two ecosystems characterized by recent population expansion. *Heredity*. doi: 10.1038/s41437-017-0019-0.

Puckett EM, Kristensen TV, Wilton CM, Lyda SB, Noyce KV, Holahan PM, Leslie DM Jr, Beringer J, Belant JL, White D Jr, Eggert LS. 2014. Influence of drift and admixture on population structure of American black bears (*Ursus americanus*) in the Central Interior Highlands, USA, 50 years after translocation. *Molecular Ecology* 23:2414–2427.

Puckett, E.E., Davis, I.S., Harper, D.C., Wakamatsu, K., Battu, G., Belant, J.L., Beyer, D.E., Carpenter, C., Crupi, A.P., Davidson, M. and DePerno, C.S., 2023. Genetic architecture and evolution of color variation in American black bears. *Current Biology*, 33(1), pp.86-97.

Sollmann R, Gardner B, Belant JL, Wilton CM, Beringer J. 2016. Habitat associations in a recolonizing, low-density black bear population. *Ecosphere* 7:8.

Tucker, M. A., Schipper, A. M., Adams, T. S., Attias, N., Avgar, T., Babić, N. L., ... & Stacy-Dawes, J. (2023). Behavioral responses of terrestrial mammals to COVID-19 lockdowns. *Science*, 380(6649), 1059-1064.

Wilton CM, Belant JL, Beringer J. 2014. Distribution of American black bear occurrences and human–bear incidents in Missouri. *Ursus* 25:53–60.

Wilton CM, Puckett EE, Beringer J, Gardner B, Eggert LS, Belant JL. 2015. Trap array configuration influences estimates and precision of black bear density and abundance. *PLOS One* 9:e111257.

Wilton CM, Beringer J, Puckett EM, Eggert LS, Belant JL. 2016. Spatiotemporal factors affecting detection of black bears during noninvasive capture-recapture surveys. *Journal of Mammalogy* 97:266–273.