


A GUIDE TO MISSOURI'S

FRESHWATER MUSSELS



MISSOURI DEPARTMENT OF CONSERVATION

A GUIDE TO MISSOURI'S
**FRESHWATER
MUSSELS**

By

Stephen E. McMurray, Caleb R. Bolin, and J. Scott Faiman,
Missouri Department of Conservation,

and

Andy Roberts & Bryan Simmons,
U.S. Fish and Wildlife Service

Photographs by
Ben Nickelson and Cliff White

Illustrations by **J. Scott Faiman**

Design by **Amanda DeGraffenreid and Marci Porter**



Front cover: Missouri has a diverse and ecologically important, but seldom seen, freshwater mussel fauna.



mdc.mo.gov

Copyright © 2012–2026 by the
Conservation Commission of the State of Missouri

Published by the Missouri Department of Conservation,
PO Box 180, Jefferson City, Missouri 65102–1080

Equal opportunity to participate in and benefit from programs of the Missouri Department of Conservation is available to all individuals without regard to their race, color, religion, national origin, sex, ancestry, age, sexual orientation, veteran status, or disability. Questions should be directed to the Department of Conservation, PO Box 180, Jefferson City, MO 65102, 573-751-4115 (voice) or 800-735-2966 (TTY), or to Chief, Public Civil Rights, U.S. Department of the Interior, 1849 C Street, NW, Washington, D.C. 20240.

TABLE OF CONTENTS

Acknowledgments	3
Introduction	4
Importance of Mussels.....	4
Mussel Reproduction	4
Habitat in General	6
Diversity and Conservation Status.....	6
Reasons for Decline	6
Invasive Bivalves.....	7
Collecting and the <i>Wildlife Code</i>	8
Observing Mussels.....	8
Reporting Discoveries.....	9
Introduction to Species Accounts	10
Status Definitions.....	10
Shell Characteristics.....	11
Mussel Shell Anatomy.....	12
Distribution Maps	13
Length Measurements	13
Species Accounts	15
Additional Species Possibly Occurring in Missouri	83
Invasive Species	87
Glossary	90
References	92
Index.....	93

ACKNOWLEDGMENTS

Bernard Sietman (Minnesota Department of Natural Resources) and Kevin Cummings (Illinois Natural History Survey) are thanked for critically reviewing and improving the previous edition of this field guide. Dyan Pursell, Jada O'Donnell, and Sarah Hobson (MDC) are thanked for assisting with data entry and the distribution maps. Alison Stodola and Rachel Vinsel (Illinois Natural History Survey), G. Thomas Watters (The Ohio State University Museum of Zoology), and John L. Harris (Arkansas State University Museum of Zoology) provided specimens for photographing.



Missouri's freshwater mussels, like the yellow sandshell (*Lampsilis teres*, page 42), live secretively on the bottom of our lakes and rivers, where they provide valuable benefits to our aquatic ecosystems. *Photo by Caleb R. Bolin.*


INTRODUCTION

Importance of Mussels

The diversity of freshwater mussels found in Missouri is an extraordinary component of our natural heritage. Living mussels are secretive and seldom seen, but the array of lustrous, ornate, and textured shells that they leave behind provides us with a familiar and appealing presence in our rivers. We are only beginning to discover the biology of mussels and the remarkable behaviors surrounding their interesting life cycle. Mussels provide vital functions in aquatic ecosystems. They are an important food source for a variety of animals, including fish, small mammals, and some birds. They filter algae, bacteria, and other particles from the water, improving water quality and cycling nutrients and energy in streams and lakes. Mussels are often used in conservation management as indicators of the overall health of aquatic ecosystems and water quality. Historically, the shells of mussels were an important resource in button manufacturing, and some species are still commercially important in the cultured pearl and jewelry industries.

Mussel Reproduction

The freshwater mussel life cycle is unique. Males release sperm, which are filtered from the water by females. The fertilized eggs are brooded in the female's gills and develop there into tiny larvae called glochidia. The glochidia are parasitic and must attach to the gills, fins, or skin of their host, usually a fish, to complete their development. Some mussel species can use a variety of species as host while others can only develop on just a single host species. The ways in which mussels attract their host and deliver their larvae are varied and fascinating. Some species, including members of the genus *Lampsilis* and its relatives, display lures that resemble small fish or crayfish to attract fish such as bass or walleye, which strike at the lure and inhale the larval mussels. Other mussels release packages of unfertilized eggs and glochidia called conglomerates, which resemble prey items like insects or worms that rupture when the fish attempts to eat them, releasing the glochidia. Glochidia are not harmful to the hosts and remain on the host for only a few days or weeks before dropping off and continuing their development into adult mussels.



- 1** Males release sperm into water. Females filter the sperm into their gill chambers where the eggs are fertilized, subsequently developing into tiny larvae called glochidia.
- 2** Glochidia lack internal organs and must attach to the gills or fins of a specific host to develop. Some mussels must trick the host into biting or inhaling the glochidia, using lures resembling small fish, insects, or other prey items of the host.
- 3** When developed, juvenile mussels separate from the host and drop to the stream bottom. Although parasitic, glochidia usually do not harm their host.
- 4** If the juvenile mussels drop into good habitat their chances for survival are much greater, enabling them to reach adulthood and reproduce.

Habitat in General

Mussels can potentially be found in any permanent body of water in Missouri. A few species can be found in ponds, lakes, and reservoirs. However, most species require flowing water and live only in streams and rivers. Mussel diversity and abundance increases downstream, as the river is joined by tributaries and increases in size. Adult mussels live partly buried in the substrate. Although some species actively burrow and move around, others may remain in one spot for decades. Most river-dwelling mussels have similar habitat requirements, and as many as 20 or more species can be found living together in favorable areas, called mussel beds. Mussel beds form in the most stable areas in streams, where the substrate is usually a firm mixture of cobble, gravel, sand, and silt. Stable areas are solid underfoot, and deep enough to be submerged even during drought. Unstable areas generally have few mussels because the loose gravel and sand move during floods, displacing or burying the mussels.

Diversity and Conservation Status

Approximately 67 species of native mussels have been found in Missouri, and at least two others may occur here. The exact number is not known because discoveries are still being made and the nomenclature of freshwater mussels fluctuates. Species composition varies by river system, and Missouri's mussel fauna is diverse because several major river systems with distinctive faunas are present. Some species have narrow ranges and are restricted to a few rivers (and some to just a few locations). Other species are widespread and may be found throughout the state. Mussels reach their highest diversity in Missouri in streams that drain the Ozark Uplift, including the Meramec, Gasconade, Osage, Spring, White, Black, and St. Francis rivers. Nearly 45 percent (30 species) of the mussel species in Missouri are species of conservation concern. Of these, 16 are in danger of extirpation in Missouri or of extinction throughout their range. For several endangered species, Missouri rivers are home to some of the last remaining populations of these rare animals.

Reasons for Decline

Freshwater mussels are sensitive to water quality degradation and alteration of physical habitat. The construction of dams to form reservoirs has been a major factor in the decline of native mussels. Reservoirs eliminate long stretches of flowing water that are vital for the survival of most mussel species. Downstream of dams, flow,

temperature, water chemistry, and fish host populations are altered. Dredging, channelization, gravel mining, and removal of streamside trees can destabilize streams for decades, seriously impacting mussel recruitment. Sediments entering streams from cleared land can bury mussels. Mussels are also sensitive to many types of pollution including metals, ammonia, and pesticides. Nonnative fish, such as invasive carp, round goby, or snakehead, can displace native fish hosts or be predators of mussels, and the invasive zebra mussel and Chinese basket clam compete with native mussels for food and space. The conservation of native mussels in the future will depend upon good stewardship of watersheds that will benefit all aquatic life.

Invasive Bivalves

Invasive bivalves, including basket clams of the genus *Corbicula* (Chinese basket clam, *Corbicula fluminea*, and Yangtze basket clam, *Corbicula largillierti*, and an undescribed species) and the zebra and quagga mussels (*Dreissena polymorpha* and *D. bugensis*), have contributed to the decline of native mussel populations. These invasive species do not have the complex life cycle of native mussels and are more closely related to the native pea clams. The Chinese basket clam was first discovered in the United States in the late 1930s, and Missouri in 1961. It quickly spread across the continent and is now nearly ubiquitous, where it occupies both rivers and lakes, and can reach densities up to 10,000 per square meter. It is believed to compete with native mussels for food and space, and large die-offs of Chinese basket clam can poison native mussels by increasing ammonia levels.

The introduction of the *Dreissena* species to North America in the 1980s resulted from ocean-crossing vessels from Europe that discharged freshwater ballast containing the free-swimming microscopic larvae into the Great Lakes. The larvae and adults have since been continually spread into new areas by humans as well as by drifting downstream. Adult *Dreissena* attach themselves to virtually any hard surface, including boat hulls, outboards, gear, and even aquatic vegetation, and can be easily transported when boats, gear, and water are moved from place to place. Zebra mussels were first detected in Missouri in 1992 and are present in several Missouri reservoirs, including Lake of the Ozarks, Truman Lake, Bull Shoals Lake, Lake Taneycomo, Lake Lotawana, Smithville Lake, Prairie Lee Lake, Lake Jacomo, Blue Springs Lake, Carrollton Recreation Lake, and Longview Lake, and several rivers, including the Osage River, Missouri River, Mississippi River, and the lower Meramec River. Zebra

and quagga mussels starve and suffocate native mussels by attaching to their shells and the surrounding habitat in large numbers. The spread of these prolific species has caused severe declines of native freshwater mussel species in many areas. The spread of zebra mussels can be thwarted by remembering to clean, drain, and dry boats and other gear that is used in the water, and by disposing of unused bait in the trash.

Collecting and the *Wildlife Code of Missouri*

In Missouri, it is legal to collect most mussel species with a Resident or Nonresident Fishing Permit. There is a daily limit of five live or dead mussels, with two shell halves counting as one animal. Chinese basket clams may be taken and possessed in any number. Zebra mussels and quagga mussels are prohibited species in Missouri and may not be collected or possessed. Mussels may be collected year-round by hand, hand net, or pole and line, and can be used as bait. Written permission from the MDC director is required to collect Missouri endangered species and species of conservation concern, so don't collect animals unless you are sure of their identification. A federal endangered species permit is required to collect species listed as endangered or threatened by the U.S. Fish and Wildlife Service. A Wildlife Collector's Permit is required to collect or possess mussels for scientific or educational purposes. Other restrictions may apply in certain areas of the state. For MDC areas, *always check* the area regulations before doing any collecting. Other agencies and organizations, including The Nature Conservancy, Missouri Department of Natural Resources, U.S. Forest Service, and National Park Service, also regulate collecting on their lands. Finally, remember that much of the land in Missouri is privately owned. Always ask for permission before entering private property.

Observing Mussels

Dead shells can be found by walking banks during low water periods. Live animals can be located by snorkeling, using water scopes, wading or hand-groping in shallow water, and SCUBA or surface-supplied air in deeper waters (appropriate safety certifications are required). For your activity to have the most value, carry a camera and a field journal to take notes on the exact location, the habitats and species present, the condition of the animals (live or dead shells), and sketch

maps to show locations and indicate stream conditions. Good digital photographs can be nearly as useful as shells for documenting your finds, and they are much easier to store and share. Records of rare species, mussel die-offs, or new localities for native or invasive species are of interest to the professional community and should be reported. Finally, return all live animals to the stream as close to where they were found as possible. To return a mussel to the stream, make sure that the hinge where the two halves of the shell are joined is pointing up.

Reporting Discoveries

If you think that you have discovered something significant about mussels, such as a diverse mussel bed, a native or invasive species outside its known range, a rare species, or a recent die-off of mussels (many dead animals with tissue still in the shells), please report it. To document species identification, please do the following:

1. If the mussel is live, take numerous high-quality photos (in the shade if possible) of both sides of the animal. Close-ups of the umbos ("beaks") can be very useful if they are not worn away.
2. Return live animals to the river near their original location. You can partly embed the mussel, hinge side up, or lay it flat on the substrate.
3. Up to five dead animals (10 shell halves) can be collected (See Collecting and the *Wildlife Code of Missouri*). Place them in a labeled bag with river name, nearest bridge crossing, or other available location information (i.e. township, range, section or GPS coordinates), collector(s) name, date, and suspected species name.
4. Take wide angle photos of the immediate vicinity to help indicate where the mussel was collected.
5. Sketch a map of the river location, indicating where the specimen was located and/or returned, where photos were taken, and any other unique features to help identify the site. Include your put-in and take-out locations and landowner contact information if permission was granted.
6. Report your findings to your nearest MDC regional office.

INTRODUCTION TO SPECIES ACCOUNTS

Two freshwater mussel families occur in Missouri — Margaritiferidae and Unionidae. Spectaclecase (*Cumberlandia monodonta*) is our only species in the family Margaritiferidae; all other native species belong to the family Unionidae. The Unionidae are further divided into tribes — a level between family and genus — and there are five tribes represented in Missouri: Anodontini, Amblemini, Lampsilini, Pleurobemini, and Quadrulini. Species accounts are arranged by family, by tribe, and then alphabetically by genus and species. Each account includes the species status in Missouri, the most distinguishing shell characteristics (including those used to separate similar looking species), Missouri distribution, habitat, and similar species for comparison.

Status Definitions

Federal endangered indicates species that are recognized by the U.S. Fish and Wildlife Service to be in danger of extinction throughout all or a significant portion of their range.

Federal threatened species are recognized by the U.S. Fish and Wildlife Service as likely to become endangered within the foreseeable future.

Species of conservation concern are species considered to be critically imperiled (S1), imperiled (S2), or vulnerable (S3) in Missouri.

State endangered species are those animals listed in Rule 3 CSR 10-4.111 of the *Wildlife Code of Missouri* as endangered in Missouri.

Common species are those that can be encountered with relative ease in the proper habitat and may be widespread in the state.

Common, but restricted indicates those species that are common, but are restricted to a particular part of the state or river drainage.

Undetermined species are those that are known to occur in Missouri, but there isn't enough information available to determine their status.

Unknown indicates species that have been reported in Missouri, but we are uncertain of their occurrence.

Invasive species are those that are not native to Missouri.


Keep in mind that the status of species may change. You should stay current with the status of Missouri's mussel species before collecting any animals. Updated annually, the Missouri Species and Communities of Conservation Concern Checklist (available at mdc.mo.gov) provides the status of Missouri's species of conservation concern.

Shell Characteristics


The shell characteristics listed for each species are those that can be used to identify and separate them from other, similar species. Refer to the shell shape image, shell characteristics image and glossary for unfamiliar terms. Keep in mind that freshwater mussels are highly variable in size and shape, and even experts can have difficulty identifying some individuals to species. The images included in this field guide are representative of the species in Missouri, but don't expect any individual to match the pictures exactly. Learning to identify mussels requires looking at a large number of individuals to learn the range of variation within each species. One of the most important variables is age. Like people, mussels change shape as they get old. A very useful trick is to look at the early growth lines of old shells. In that way you can see what the shape of the shell was when it was younger.




Rhomboidal




Round



Oval



Oval



Quadrat



Elliptical



Triangular





Concave Posterior Ridge




Convex Posterior Ridge

Mussel Shell Anatomy



Distribution Maps


The distribution map included with each species account is based on known occurrences at the time of publication and represent the occurrence of live animals or shell material, from collections, published sources, or museum records. Our knowledge is incomplete, and you might very well find a new location for some of our mussel species! The maps are only intended to show the potential distribution of each species in Missouri and should not be used to infer conservation status. Some species are rarely encountered, and even though they may be known from several river basins in the state, it might literally take experienced collectors an entire day to even find an old shell.



River drainages highlighted in blue indicate where a live mussel or shell material has been observed.

Length Measurements

The scale bar included with each account shows the length of the actual specimen pictured. Also included is the maximum recorded size for each species.




SPECIES ACCOUNTS

Spectaclecase

Cumberlandia monodonta (Say, 1829)

Status: Federal Endangered
State Endangered
Species of Conservation Concern



Shell characteristics: Shell thin to moderately thick; elongate and compressed, often arched in shape; smooth. Posterior ridge rounded, long; shell turned down posteriorly. Pseudocardinal teeth poorly developed, conical, lateral teeth nearly absent; no beak cavity, beak sculpture of two to three parallel lines. Periostracum black or brown, often peeling; rays absent. Nacre bluish white to white.

Habitat: Large rivers, in gravel and cobble, often in association with boulders or under large, flat rocks, and bluff pool habitats. The Meramec and Gasconade rivers may have the largest populations of spectaclecase in the United States.

Similar species: spike, black sandshell



5.7"


Length: Up to 10"

Elktoe

Alasmodonta marginata Say, 1818

Status: Species of Conservation Concern

Shell characteristics: Shell thin; nearly rhomboidal to elongated, truncated posteriorly, inflated; umbos located more posterior than most mussels and elevated above the hinge line, beak sculpture of three to four strong, rhomboidal, wavelike, concentric ridges. Posterior slope broad and flat, sculptured with low, oblique ribs, with a sharp posterior ridge. Lateral teeth absent, small pseudocardinal teeth. Periostracum yellowish to light brownish, with greenish tinge; numerous fine to wide zigzag blue-green rays, usually speckled with small dark flecks. Nacre white or iridescent.



Habitat: Creeks to large rivers, and ditches, usually found in gravel or gravel-sand substrates.

Similar species: snuffbox, slippershell mussel, deerto



2.4"

Length: Up to 4"

Cylindrical papershell

Anodontoides ferussacianus


(Lea, 1834)

Status: Species of Conservation Concern

Shell characteristics: Shell thin, fragile; elliptical, usually with a straight dorsal margin; smooth; beaks slightly inflated with two to three curved, inconspicuous concentric ridges. Posterior ridge indistinct. Both pseudocardinal and lateral teeth absent; beak cavity absent. Periostracum light green to yellowish-brown, green rays present in younger shells. Nacre silvery to bluish white.

Habitat: Can be found in small streams and in rivers in silt, gravel, or sand.

Similar species: creeper, paper pondshell, giant floater




2.4" Length: Up to 4"

Rock pocketbook

Arcidens confragosus (Say, 1829)

Status: Species of Conservation Concern

Shell characteristics: Shell thin to moderately solid; nearly quadrate to rhomboidal; inflated; beak sculpture of two prominent rows of tubercles extending across umbo. Irregular folds or ridges on posterior third of shell. Posterior ridge indistinct. Pseudocardinal teeth sharp, distinct, lateral teeth absent except for a single flange extending halfway along hinge line. Periostracum greenish to black, no rays. Nacre bluish white.



Habitat: Ditches, sloughs, creeks, and medium-to-large rivers in mud in slow current.

Similar species: threeridge, washboard



3.9"


Length: Up to 6"

Flutedshell

Lasmigona costata (Rafinesque, 1820)

Status: Common, but restricted

Shell characteristics: Shell thin to moderately thick; nearly rhomboidal, compressed; beak sculpture of three to four coarse ridges, slightly notched in the middle. Posterior ridge indistinct; distinct flutings on posterior ridge in most populations. Pseudocardinal teeth weak to moderate, lateral teeth poorly developed. Periostracum yellowish-brown to black, sometimes with green rays. Nacre white.



Habitat: Found in creeks to medium-sized rivers in moderate current; prefers gravel substrates.

Similar species: creeper




3.2" Length: Up to 6"

White heelsplitter

Pterosyna complanata (Barnes, 1823)

Status: Common

Shell characteristics: Shell thin to thick, sturdy; nearly quadrate to rounded; laterally compressed, with a pronounced wing; umbos flat, beak sculpture of heavy W-shaped ridges. Usually has flutings on the wing, otherwise shell smooth. Posterior ridge indistinct. Pseudocardinal teeth moderate to weak; lateral teeth poorly developed. Dark periostracum, rayless. Nacre white.



Habitat: Found in ditches, creeks, and medium-to-large rivers with mud or mud-gravel substrates in slow current.

Similar species: pink heelsplitter, fragile papershell, pink papershell



4.5" Length: Up to 10"

Giant floater


Pyganodon grandis (Say, 1829)

Status: Common

Shell characteristics: Shell thin, fragile; nearly oval; smooth; umbos broad, moderately inflated, sculpture of three to four double-loops, almost W-shaped, sometimes asymmetrical or broken. Posterior ridge indistinct. Teeth absent. Periostracum rayless and usually brown but can be yellow or green. Nacre iridescent or salmon to white.

Habitat: Prefers slack water habitats, with mud or gravel bottoms, in ditches and small creeks to large rivers; can be found in lakes or ponds.

Similar species: creeper, cylindrical papershell, flat floater, pondhorn




4.5" Length: Up to 12"

Salamander mussel

Simpsonaias ambigua (Say, 1825)

Status: Proposed Federal Endangered
State Endangered
Species of Conservation Concern



Shell characteristics: Shell thin; elliptical to oblong, with parallel dorsal and ventral margins, moderately compressed; smooth; beak sculpture of several thin ridges forming an upside down "V." Posterior ridge indistinct. Lateral teeth absent, pseudocardinal teeth very reduced. Periostracum brownish and rayless. Nacre dull whitish-purplish, salmon in beak cavity.

Habitat: Found under large, flat rocks, often in mud, gravel or cobble, in medium-sized rivers; has also been reported living inside of dead shells of larger species.

Similar species: cylindrical papershell, spectaclecase, pondhorn



1.3" Length: Up to 2"

Creeper

Strophitus undulatus (Say, 1817)


Status: Common

Shell characteristics: Shell thin, strong; quadrate to elliptical, compressed to moderately inflated; smooth; beaks slightly elevated above the hinge line and located near the middle of the shell, sculpture of two to three coarse ridges. Anterior end rounded, posterior end bluntly pointed.

Posterior ridge indistinct. Lateral teeth absent, pseudocardinal teeth poorly developed. Lateral-pseudocardinal teeth form a distinctive waved outline. Periostracum green with rays in juveniles, becoming chestnut, dark brown, or black in older individuals, shiny. Nacre iridescent, salmon to milky white in umbo.

Habitat: Found in small-to-large streams, preferring gravel or gravel-mud substrates.

Similar species: cylindrical papershell, giant floater, paper pondshell



2.7"


Length: Up to 4"

Slippershell mussel

Sulcularia viridis (Rafinesque, 1820)

Status: State Endangered
Species of Conservation Concern

Shell characteristics: Shell moderately thick; rectangular to nearly trapezoidal; often inflated, with a high posterior ridge; smooth; beak sculpture composed of several heavy, concentric loops. Posterior slope abrupt, smooth. Lateral teeth absent, pseudocardinal teeth triangular and distinct. Periostracum is often a semi-metallic silvery-gray or yellowish with numerous green or brownish wavy rays. Nacre silvery white to iridescent.



Habitat: Found in small streams to medium-sized rivers in sand, mud, or fine gravel, often near water willow.

Similar species: elktoe, snuffbox




1.5" Length: Up to 2"

Paper pondshell

Utterbackia imbecillis (Say, 1829)

Status: Common

Shell characteristics: Shell thin, fragile; nearly elliptical, inflated; hinge line nearly straight; smooth; umbos flattened and do not rise above the hinge line, sculptured with flat double loops. Posterior ridge indistinct. Lateral and pseudocardinal teeth absent. Periostracum green or yellowish green, often with fine green rays. Nacre white to bluish.



Habitat: Ditches and small creeks to large rivers, prefers backwaters, sloughs, ponds, and oxbows, usually in silt or silt and sand.

Similar species: cylindrical papershell, giant floater, scaleshell



2.2"


Length: Up to 4"

Flat floater

Utterbackiana suborbiculata (Say, 1831)

Status: Species of Conservation Concern

Shell characteristics: Shell thin, fragile; round, nearly circular; smooth; compressed with a low dorsal keel; hinge line straight. Posterior ridge indistinct. Teeth absent. Beak sculpture composed of two rows of broken ridges appearing double looped. Periostracum yellow, tan, and green with very faint color rays, especially in young specimens. Nacre white to pinkish, iridescent.



Habitat: Ditches and creeks to large rivers in silt. Can be locally abundant, especially in oxbow lakes and backwaters of large rivers.

Similar species: giant floater, paper pondshell




3.8" Length: Up to 12"

Threeridge

Amblema plicata (Say, 1817)

Status: Common

Shell characteristics: Shell moderately thick, solid; quadrate to rounded, moderately inflated to inflated; beak sculpture of two to three curved concentric ridges. Posterior ridge present, shell with folds or undulations parallel to the posterior ridge, mainly posterior to umbo. Stout and heavy pseudocardinal and lateral teeth. Periostracum brownish to black. Nacre white, sometimes iridescent blue at posterior end ("bluepoint").



Habitat: Creeks, ditches, small-to-large rivers, lakes, in gravel, sand, or mud.

Similar species: washboard, bankclimber, rock pocketbook



3.4"


Length: Up to 6"

Western rainbow

Cambarunio hesperus Watters, 2018

Status: Common, but restricted

Shell characteristics: Shell thin, especially on posterior tip, but strong; elongate to elliptical, compressed; small; umbos small, only slightly elevated, sculpture of several ridges, 2–3 concentric, others appearing double-looped. Posterior ridge indistinct, shell smooth. Pseudocardinal teeth triangular, lateral teeth thin and bladelike; beak cavity shallow. Periostracum yellow to yellow-green, with distinctive broad green rays on posterior third of shell. Nacre iridescent to white or bluish-white.



Habitat: Creeks and small-to-medium rivers in gravel.

Similar species: ellipse, pondmussel, Texas lilliput, little spectaclecase, bleedingtooth mussel




1.5" Length: Up to 3"

Western fanshell

Cyprogenia aberti (Conrad, 1850)

Status: Federal Threatened
State Endangered
Species of Conservation Concern



Shell characteristics: Shell thick, strong; round to triangular, compressed to moderately inflated; umbos low, sculpture faint. Posterior ridge prominent and raised, shallow sulcus from umbo to middle of the ventral margin; shell with a wrinkled or rough appearance. Pseudocardinal teeth large, lateral teeth short and slightly curved; beak cavity moderately deep. Periostracum dull, tan, with distinctive ray pattern from bands of tiny pigment flecks. Nacre white.

Habitat: Creeks and small-to-medium rivers in gravel.

Similar species: pimpleback, deertoe




— 2.0" — **Length:** Up to 3"

Butterfly

Ellipsaria lineolata (Rafinesque, 1820)

Status: Common, but restricted

Shell characteristics: Shell stout, thick; triangular and compressed; umbos broad, flattened, pointed at the apex, directed forward, sculpture of multiple fine double-looped lines. Posterior ridge prominent and sharply angled; sexually dimorphic — females are smaller and more robust (thickened posteriorly) than males. Lateral and pseudocardinal teeth large and well defined; beak cavity shallow to moderately deep. Periostracum yellowish to brownish, becoming darker with age, with characteristic dark brown raying. Nacre white.



Habitat: Creeks, small-to-large rivers in stable gravel.

Similar species: deertoe, mucket



3.0"


Length: Up to 4"

Curtis pearlymussel

Epioblasma curtisii

(Frierson and Utterback, 1916)


Status: Federal Endangered
State Endangered
Species of Conservation Concern



Shell Characteristics: Shell relatively thick, stout; oval to rhomboidal, inflated; umbos broad, raised slightly above the hinge line, sculpture unknown. Posterior ridge of male biangulate, of female broadly rounded; shell with prominent growth lines; sexually dimorphic — females with distinct marsupial swelling and more rhomboidal in outline. Pseudocardinal and lateral teeth well defined, beak cavity broad. Periostracum yellow-tan with green rays. Nacre white.

Habitat: Small Ozark streams to large rivers in stable gravel.

Similar species: snuffbox, ellipse



Snuffbox

Epioblasma triquetra



(Rafinesque, 1820)

Status: Federal Endangered
State Endangered
Species of Conservation Concern

Shell characteristics: Shell thick and stout; oblong to oval, inflated; umbos broad, slightly raised, sculpture of faint, double-looped ridges. Posterior ridge strong, concave, with fine parallel ribbing; sexually dimorphic — females with the posterior part of the shell laterally expanded (looks almost “nose-like”); this expansion is finely ribbed, the ends of the ribs forming sharp teeth on the ventral margin. Pseudocardinal teeth elevated, thin, and rough; lateral teeth short and curved; beak cavity deep. Periostracum yellowish or greenish, with bold interrupted rays, occasionally forming chevron markings. Nacre pearly white.

Habitat: Small-to-medium rivers in gravel.


Similar species: elktoe, deertoe, fawnsfoot



Pink mucket

Lampsilis abrupta (Say, 1831)


Status: Federal Endangered
State Endangered
Species of Conservation Concern



Shell characteristics: Shell thick, heavy, solid; slightly rounded to broadly elongate, rounded or blunt posterior end; beaks low, not raised above the hinge line, sculpture of six to 10 fine, wavy double-looped ridges. Posterior ridge indistinct, shell smooth; sexually dimorphic — females are square in shape, males are round. Pseudocardinal teeth stout, triangular, erect; lateral teeth long, curved, sharp; beak cavities deep. Periostracum yellowish brown to chestnut, sometimes with faint rays (juveniles or younger individuals). Nacre white, with pinkish or salmon beak cavity; sometimes all white.

Habitat: Medium-to-large rivers in gravel, sand, cobble.


Similar species: mucket, plain pocketbook, Higgins eye



Missouri brokenray

Lampsilis britis Simpson, 1900


Status: Species of Conservation Concern
Missouri Endemic



Shell characteristics: Shell thin but strong; elliptical to ovate or elongate, compressed to slightly inflated; umbos large, but only slightly raised above the hinge line, beak sculpture of inverted, V-shaped ridges. Posterior ridge indistinct; shell smooth; sexually dimorphic — females resemble a “catcher’s mitt” in outline with an indentation on the post-ventral margin. Pseudocardinal teeth distinct, lateral teeth short, slightly curved, taller posteriorly; beak cavity shallow. Periostracum light yellowish, becoming darker with age; green rays fine to broad, more dense posteriorly, appearing splotchy. Nacre white to bluish-white, iridescent posteriorly.

Habitat: Occurs in small streams and headwaters of medium rivers in gravel substrate. Also known as northern brokenray, occurs in streams that flow north off the Salem and Springfield plateaus.

Similar species: Arkansas brokenray, fatmucket, ellipse




Plain pocketbook

Lampsilis cardium Rafinesque, 1820


Status: Common

Shell characteristics: Shell moderately thick, stout; quadrate to oval or elliptical, moderately to very inflated; beaks broad, raised above the hinge line, sculpture of two to three coarse ridges. Posterior ridge indistinct and rounded; shell smooth; sexually dimorphic — posterior end sharp in males, broadly rounded in females. Pseudocardinal and lateral teeth distinct, solid; beak cavities broad and deep. Periostracum usually yellow, sometimes tan or brown, thin rays usually present. Nacre white, often pinkish in beak cavity.



Habitat: Small creeks to large rivers, in a variety of substrates.


Similar species: mucket, pink mucket, bleufer, fat pocketbook, fatmucket



Higgins eye

Lampsilis higginsii (Lea, 1857)


Status: Federal Endangered
State Endangered
Species of Conservation Concern



Shell characteristics: Shell thick, heavy; rounded to slightly elongate, moderately inflated; beaks broad, above the hinge line, directed forward, sculpture of fine double-looped lines. Posterior ridge indistinct, rounded; shell smooth; sexually dimorphic — females rounded and truncate, males oval. Large, triangular pseudocardinal teeth; lateral teeth long, heavy, straight; beak cavity deep. Periostacum yellowish-brown, with bold green rays. Nacre white, sometimes tinged with salmon or cream.

Habitat: Large rivers in gravel or sand; currently only known from the mainstem upper Mississippi River in Missouri.

Similar species: pink mucket, mucket, plain pocketbook




2.8" Length: Up to 6"

3.5" Length: Up to 6"

Neosho mucket

Lampsilis rafinesqueana Frierson, 1927

Status: Federal Endangered
State Endangered
Species of Conservation Concern



Shell characteristics: Shell relatively thin, strong; oblong to quadrate, compressed; beaks low, only slightly raised above the hinge line, sculpture of fine lines rounded up posteriorly. Posterior ridge broadly rounded to flattened, shell appearing biangulate; shell smooth. Pseudocardinal teeth stout, lateral teeth short and stout, downcurved; beak cavity relatively shallow. Periostracum yellow to tan; shell with interrupted broad green rays or chevrons. Nacre bluish white to white, slightly iridescent posteriorly.

Habitat: Small-to-medium streams in gravel or mixture of gravel and sand substrate; occurs exclusively in streams of the Neosho River system in southwest Missouri.

Similar species: mucket, plain pocketbook, fatmucket



2.8"


Length: Up to 5"

Arkansas brokenray

Lampsilis reeveiana (Lea, 1852)


Status: Common, but restricted

Shell characteristics: Shell thin; elliptical to ovate, moderately inflated; umbos small, only slightly raised above the hinge line, beak sculpture of inverted, V-shaped ridges. Posterior ridge indistinct, broadly rounded; shell smooth, slightly alate posteriorly. Pseudocardinal teeth small but distinct, lateral teeth relatively short and curved; beak cavity shallow. Periostracum yellowish to yellowish brown, shiny; numerous thin to broad green rays present. Nacre salmon to bluish-white, iridescent posteriorly.



Habitat: Small streams and headwaters of medium streams in gravel substrate; only occurs in streams that flow south off the Salem and Springfield plateaus.

Similar: Missouri brokenray, fatmucket, bleedingtooth mussel



Canary kingshell

Lampsilis sietmani


Keogh and Simons, 2019

Status: Undetermined

Shell characteristics: Shell thick, robust; oblong and distinctly elongate, moderately inflated; umbos broad, raised above the hinge line, indistinctly sculptured or with discontinuous ridges. Posterior ridge rounded; shell smooth; sexually dimorphic – females swollen posteriorly. Pseudocardinal teeth robust and erect, lateral teeth long, straight to slightly incurved; beak cavity shallow. Periostracum shiny, straw-colored, vibrant yellow or a stained khaki, green rays only present in young individuals. Nacre white to blue-white, iridescent posteriorly.

Habitat: Medium-to-large rivers in a variety of substrates.

Similar: yellow sandshell, mucket, plain pocketbook, fatmucket




5" 4.9" Length: Up to 7"

Fatmucket

Lampsilis siliquoidea (Barnes, 1823)

Status: Common

Shell characteristics: Shell thin to thick, solid; elongate to elliptical, moderately inflated; umbos large, only slightly raised above the hinge line, sculpture of multiple fine double-looped lines. Posterior ridge not prominent; shell smooth; sexually dimorphic — females prominently swollen posteriorly. Pseudocardinal and lateral teeth thin, erect; beak cavity shallow. Periostracum typically yellowish, usually with distinct rays. Nacre bluish-white to white.



Habitat: Small streams to large rivers, lakes, reservoirs, in a variety of substrates.

Similar species: yellow sandshell, canary kingshell, mucket, plain pocketbook




2.7" 3.6" Length: Up to 6"

Yellow sandshell

Lampsilis teres (Rafinesque, 1820)

Status: Common

Shell characteristics: Shell moderately thick, solid; elongate, moderately inflated; umbos large, only slightly raised above the hinge line, sculpture of multiple fine double-looped lines. Posterior ridge indistinct; shell smooth; sexually dimorphic — females swollen posteriorly. Pseudocardinal teeth distinct and elevated, lateral teeth distinct and long; beak cavity shallow. Periostracum distinct yellow to yellowish-tan, often shiny. Nacre white.



Habitat: Small-to-large rivers, oxbows, or reservoirs in sand to fine gravel.

Similar species: black sandshell, fatmucket, canary kingshell




3.2" Length: Up to 8"

Little spectaclecase

Leaunio lienosus (Conrad, 1834)

Status: Species of Conservation Concern

Shell characteristics: Shell thin but strong; elongate to elliptical, moderately inflated; small; umbos moderately inflated and slightly elevated, sculpture of four to six fine double-looped lines. Posterior ridge indistinct; shell smooth; sexually dimorphic — females with posterior end inflated and truncated. Pseudocardinal teeth triangular, lateral teeth short and curved; beak cavity moderately deep. Periostracum brown to tan, becoming darker with age, sometimes with greenish rays or rayless. Nacre iridescent, pink, sometimes creamish or bluish-white.



Habitat: Small-to-medium streams in gravel.

Similar species: pondmussel, Texas lilliput, purple lilliput




1.7" Length: Up to 3"

Black sandshell

Ligumia recta (Lamarck, 1819)

Status: Species of Conservation Concern

Shell characteristics: Shell thick, heavy, solid; elongate, compressed to moderately inflated; umbos low, barely above the hinge line, sculpture of two to three double-looped lines, usually obscure. Posterior ridge indistinct; shell smooth; sexually dimorphic — posterior end of females is blunt and turned upwards. Pseudocardinal teeth well developed but not large, lateral teeth elongate and straight; beak cavity shallow. Periostracum polished black to dark brown with obscure green rays that are more prevalent in young. Nacre is white to iridescent, often pink to purplish.



Habitat: Medium-to-large rivers in gravel to firm sand.

Similar species: spike, yellow sandshell, canary kingshell, spectaclecase, pondmussel



5.8"


Length: Up to 8"

Threehorn wartyback

Obliquaria reflexa (Rafinesque, 1820)

Status: Common

Shell characteristics: Shell thick, stout; roundly triangular, somewhat inflated; umbos narrow, incurved, above the hinge line, sculpture of two to three ridges, more prominent posteriorly. Posterior ridge broadly rounded and prominent, with slight flutings on posterior slope. Single rows of knobs that alternate on both valves. Pseudocardinal teeth prominent, lateral teeth short and straight; beak cavity shallow. Periostracum light tan to green, with numerous interrupted fine rays. Nacre pearly white.



Habitat: Medium-to-large rivers in gravel.

Similar species: pimpleback, sheepnose, wartyback



2.3"


Length: Up to 3"

Southern hickorynut

Obovaria arkansasensis (Lea, 1862)

Status: Species of Conservation Concern

Shell characteristics: Shell thick, stout; oval to elliptical, gently rounded dorsally and ventrally, moderately inflated; umbos broad and low but raised above the hinge line, sculpture unknown. Posterior ridge not prominent; sexually dimorphic — females have a bluntly truncated posterior end. Pseudocardinal teeth well defined and triangular, lateral teeth high, roughened, and slightly curved; beak cavity shallow. Periostracum light to dark brown, with faint green capillary rays on posterior two-thirds. Nacre bluish-white, iridescent posteriorly.



Habitat: Small-to-medium rivers in sand or gravel.

Similar species: hickorynut, round pigtoe, Wabash pigtoe




Hickorynut

Obovaria olivaria (Rafinesque, 1820)

Status: Species of Conservation Concern

Shell characteristics: Shell thick, stout; oblong to ovate, inflated; umbos narrow, high, and directed forward, sculpture of a series of fine lines, often offset in the middle and drawn up. Posterior ridge broadly rounded, dorsal margin nearly straight, anterior end broadly rounded, posterior end sharply rounded. Large, thick, well-defined teeth; beak cavity shallow. Periostracum olive-brown to yellowish-brown, with green capillary rays present, some may be rayless, often a slight sheen. Nacre blue-white, iridescent throughout.



Habitat: Large rivers in mixed sand or gravel.

Similar species: southern hickorynut, round pigtoe, ebonyshell, Higgins eye, pink mucket



1.9" Length: Up to 3"


Mucket

Ortmanniana ligamentina

(Lamarck, 1819)

Status: Common

Shell characteristics: Shell thick, moderately heavy to very heavy; oblong oval to quadrate, moderately inflated; umbos broad, only slightly elevated above the hinge line, sculpture of many fine concentric lines, often obscure. Posterior ridge broadly rounded, shell lacks any sculpturing except growth lines. Pseudocardinal teeth large, erect and somewhat triangular; lateral teeth short, downcurved, stout; relatively shallow beak cavity. Periostracum yellowish-brown, dark green rays are variable, may be broad and straight or thin, some are rayless. Nacre white, sometimes pinkish or light orange.



Habitat: Creeks to medium-to-large rivers in stable gravel.

Similar species: pink mucket, plain pocketbook, sheepnose, fatmucket



3.7" Length: Up to 7"


Bankclimber

Plectomerus dombeyanus

(Valenciennes, 1827)

Status: Species of Conservation Concern

Shell characteristics: Shell not particularly thick, but heavy; rhomboidal to quadrate, moderately inflated to compressed; umbos low, only slightly above the hinge line, sculpture of irregular double-looped ridges. Posterior ridge defined and sharp, with plications anterior to the ridge and flutings dorsal to the posterior ridge. Pseudocardinal teeth prominent, lateral teeth long and curved; beak cavity shallow. Periostracum greenish to brown, darker with age. Nacre color variable, usually purple but often tinged with bronze.



Habitat: Medium-sized streams or ditches in mud or mud-gravel; restricted to rivers, streams or ditches in southeast Missouri.

Similar species: threeridge, washboard, pistolgrip, flutedshell




5.7" Length: Up to 6"

Pink heelsplitter

Potamilus alatus (Say, 1817)

Status: Common

Shell characteristics: Shell thin but strong; quadrate, compressed; umbos low, narrow, not raised above the hinge line, sculpture of a few very fine concentric lines drawn up at either end. Posterior ridge broadly rounded; shell sculpture-less except for growth lines, with a high, prominent posterior wing; sexually dimorphic — posterior of females usually angled and bluntly squared or truncated. Pseudocardinal teeth erect, knobby, lateral teeth short, curved; beak cavity shallow. Periostracum greenish in juveniles; older individuals dark or brown, green rays may be present. Nacre uniformly purple or pink, sometimes tinged with copper.



Habitat: Creeks and medium-to-large rivers in a variety of substrates.

Similar species: white heelsplitter, pink papershell, bleufer




6.0" Length: Up to 8"

Fat pocketbook

Potamilus capax (Green, 1832)

Status: Federal Endangered
State Endangered
Species of Conservation Concern



Shell characteristics: Shell thin but stout; round to oblong, very inflated; umbos large, prominent, and turned inward, sculpture of a few faint oblique ridges. Posterior ridge sharply angled; hinge line distinctly S-shaped (most easily seen on the inside of the shell); shell has minute anterior and posterior wings. Pseudocardinal teeth erect, lateral teeth short, erect, and curved; beak cavity deep and wide. Periostracum smooth, shiny, yellow to brownish and lacking rays, highly polished. Nacre white or flushed with pink or salmon.

Habitat: Ditches, large rivers in slow flowing water, in mud or sand.

Similar species: plain pocketbook, bleufer



3.7"


Length: Up to 6"

Fragile papershell

Potamilus fragilis (Rafinesque, 1820)

Status: Common

Shell characteristics: Shell thin, fragile; oblong to ovate, compressed; umbos low and forward, sculpture of three to four delicate lines, the first concentric and remaining ones double-looped, often imperceptible. Posterior ridge broadly rounded; shell with a moderate dorsal wing (may be broken off). Pseudocardinal and lateral teeth thin, sharp; beak cavity shallow. Yellow periostracum often with thin greenish rays, or rayless. Nacre pinkish to white and highly iridescent.



Habitat: Small streams to large rivers in stable mud, sand, or gravel.

Similar species: pink papershell, scaleshell




3.5" Length: Up to 8"

Scaleshell

Potamilus leptodon (Rafinesque, 1820)

Status: Federal Endangered
State Endangered
Species of Conservation Concern



Shell characteristics: Shell thin, extremely fragile; oblong, rhomboidal to oval and compressed; umbos low and narrow, sculpture of a few very faint, double-looped ridges. Posterior ridge not prominent, shell with a weak dorsal wing; shell appears “bent” when examined from the top; sexually dimorphic — females are much smaller than males and are more elongate, and shell is flaky at posterior tip due to the periostracum layer extending past the margin of the shell. Pseudocardinal teeth poorly developed, lateral teeth weakly defined; beak cavity shallow. Periostracum olive to brown, covered with fine, wavy rays. Nacre rose colored in the cavity, bluish or purple elsewhere, iridescent throughout.

Habitat: Medium-to-large rivers in mixture of sand and gravel, also in mud.

Similar species: fragile papershell, paper pondshell




Pink papershell

Potamilus ohiensis (Rafinesque, 1820)

Status: Common

Shell characteristics: Shell very thin and fragile; oblong to oval, compressed; umbos low, sculpture of several small broken ridges with nodules, often indiscernible. Posterior ridge indistinct; shell alate, with a high, pronounced posterior wing in addition to the anterior wing, but these may be broken off in older individuals. Pseudocardinal and lateral teeth present, thin, weak. Periostracum brown to tan, shiny, some rays may be present. Nacre iridescent, pink to light purple.



Habitat: Medium-to-large rivers in a variety of substrates.

Similar species: fragile papershell, pink heelsplitter, white heelsplitter




4.4" Length: Up to 6"

Bleufer

Potamilus purpuratus (Lamarck, 1819)

Status: Common, but restricted

Shell characteristics: Shell moderately thick, stout; oval to quadrate, inflated; umbos narrow and incurved, barely above hinge line, sculpture of faint corrugations. Posterior ridge not prominent; shell slightly alate; sexually dimorphic — female posterior end usually angled and bluntly squared or truncated. Pseudocardinal teeth erect and emarginated, lateral teeth short, curved, raised; beak cavity broad and moderately deep. Periostracum dark brown to black, without rays. Nacre deep purple and highly iridescent.



Habitat: Creeks and ditches to large rivers in mud or gravel.

Similar species: pink heelsplitter, plain pocketbook



3.2"

Length: Up to 8"


Ouachita kidneyshell

Ptychobranthus occidentalis

(Conrad, 1836)

Status: Species of Conservation Concern

Shell characteristics: Shell relatively thin but stout; oblong to elongate, noticeably "kidney shaped," compressed; umbos low, sculpture of several finely spaced, inverted V-shaped ridges, often eroded. Posterior ridge not prominent; often have a finger-shaped sulcus inside the shell pointing to the umbo. Pseudocardinal and lateral teeth well developed; lateral teeth short, broad, and striated; beak cavity shallow. Periostracum yellow to brown with a cloth-like texture, may have very fine green or dark rays. Nacre white.



Habitat: Small streams in gravel. Specimens from the White and Neosho river basins may represent separate species. May also occur in the Osage, Meramec, and Gasconade river basins.

Similar species: spike, ellipse, bleedingtooth mussel




2.7" Length: Up to 5"

Pondmussel

Sagittunio subrostratus (Say, 1831)

Status: Common

Shell characteristics: Shell thin to moderately solid; elongate to elliptical, drawn to a point posteriorly above the dorsal-ventral midline, moderately inflated; small; umbos full, slightly elevated, sculptured with fine double loops. Posterior ridge indistinct; shell smooth; sexually dimorphic — females noticeably truncate. Pseudocardinal and lateral teeth distinct, thin; beak cavity shallow. Periostracum green or black with broad rays. Nacre white.



Habitat: Small creeks, ponds, lakes, in mud.

Similar species: black sandshell, little spectaclecase



Purple lilliput

Toxolasma lividum Rafinesque, 1831


Status: Species of Conservation Concern

Shell characteristics: Shell thick, stout; quadrate, with straight dorsal and ventral lines, moderately inflated; small; umbos broad, low, sculpture of three to four concentric bars, often incomplete. Posterior ridge indistinct; sexually dimorphic — females truncated posteriorly.

Pseudocardinal teeth well developed, serrated and elevated, lateral teeth straight to slightly curved; beak cavity moderately deep. Periostracum dark brown, often faint green on new parts. Nacre rusty purple fading to lighter purple or white at shell margin.

Habitat: Small-to-medium streams, lakes, ponds, in gravel, mud.

Similar species: lilliput, Texas lilliput, little spectaclecase




1.0" Length: Up to 2"

Lilliput

Toxolasma parvum (Barnes, 1823)

Status: Common

Shell characteristics: Shell moderately thin but stout; elliptical, with dorsal and ventral margins parallel, inflated; small; umbos low, broad and flattened, sculpture of five to six coarse, concentric ridges. Posterior ridge indistinct. Pseudocardinal and lateral teeth well defined but delicate, lateral teeth slightly curved; beak cavity shallow. Periostacum tan to dark brown with a cloth-like texture, posterior half tinged with green. Nacre bluish-white, iridescent throughout.



Habitat: Small-to-medium streams, lakes, ponds, in gravel, mud.

Similar species: purple lilliput, salamander mussel, Texas lilliput




1.1" Length: Up to 2"

Texas lilliput

Toxolasma texasiense (Lea, 1857)

Status: Species of Conservation Concern

Shell characteristics: Shell thin to solid; elongate, moderately inflated; umbos low, only slightly raised above the hinge line, sculpture of five to six sharp, angular ridges. Posterior ridge indistinct; sexually dimorphic — females are noticeably truncated on the posterior end and shorter in length. Pseudocardinal teeth thin and compressed, lateral teeth long, straight or curved; beak cavity shallow. Periostracum dark or greenish brown with a cloth-like texture (faint, silvery sheen). Nacre white, sometimes tinged with salmon in the beak cavity.



Habitat: Small streams to medium rivers in sand or mud; restricted to southern Missouri.

Similar species: lilliput, purple lilliput, little spectaclecase, pondmussel




1.1" Length: Up to 3"

Fawnsfoot

Truncilla donaciformis (Lea, 1828)

Status: Common

Shell characteristics: Shell relatively thick and stout; broadly triangular with a pointed posterior end, moderately inflated; umbos narrow and flattened, only slightly above hinge line, sculpture of five to six double-looped bars. Posterior ridge sharply angled. Pseudocardinal and lateral teeth moderately developed and sharp. Periostracum yellow or greenish, covered with chevron-shaped or zigzag lines. Nacre pearly white.



Habitat: Medium streams to large rivers in sand or gravel.

Similar species: deertoe




0.9" Length: Up to 2"

Deertoe

Truncilla truncata (Rafinesque, 1820)

Status: Common

Shell characteristics: Shell moderately thick and stout; triangular with posterior end bluntly pointed or truncated, inflated; umbos narrow, raised above the hinge line, sculpture of three to five thin, double-looped ridges. Posterior ridge sharply angled and prominent; sexually dimorphic – females typically have a less defined posterior ridge, straight to slightly convex ventral margin and no sulcus. Pseudocardinal and lateral teeth moderately developed. Periostracum yellow-brown to brown, often with green capillary rays grouped together to form broad bands of color, some valves may be almost rayless. Nacre white, occasionally pink.



Habitat: Medium streams to large rivers in mud, gravel, or sand.

Similar species: fawnsfoot, Wabash pigtoe, snuffbox



1.8"

Length: Up to 2"

Ellipse

Venustaconcha ellipsiformis


(Conrad, 1836)

Status: Common

Shell characteristics: Shell moderately thick, solid; elliptical to slightly ovate, slightly inflated; umbos low, sculpture of three to four fine, double-looped ridges. Posterior ridge indistinct; posterior ventral margin indented; sexually dimorphic — females have a slight crease or sulcus that runs parallel to the rays on the posterior end of the shell. Pseudocardinal and lateral teeth well developed and stout, often reddish in fresh shells. Periostracum greenish to brownish with numerous fine green rays or lines. Nacre variable, usually white, iridescent.

Habitat: Small-to-medium streams and rivers in stable gravel; common in some north-flowing Ozark streams.

Similar species: western rainbow, bleedingtooth mussel, mucket, Ouachita kidneyshell



2.2"


Length: Up to 3"

Bleedingtooth mussel

Venustaconcha pleasii (Marsh, 1891)

Status: Common, but restricted

Shell characteristics: Shell moderately thick, stout (thicker anteriorly); sub-elliptical and elongated, compressed to moderately inflated; umbos low, sculpture of three to four fine undulations. Posterior ridge broadly rounded; shell distinctly "pinched" near posterior end; sexually dimorphic — females have a pronounced sulcus running parallel to the rays on the posterior end of the shell. Pseudocardinal teeth stout and erect, lateral teeth long and nearly straight, with reddish coloring. Periostracum yellowish-brown to chestnut, with numerous fine, wavy green rays. Nacre white, center often salmon- coppery colored, iridescent.



Habitat: Small-to-medium streams and rivers in stable gravel; restricted to streams that flow south and west off the Springfield and Salem plateaus.

Similar species: ellipse, western rainbow, Ouachita kidneyshell



1.6" Length: Up to 3"

Elephantear


Elliptio crassidens (Lamarck, 1819)

Status: State Endangered
Species of Conservation Concern

Shell characteristics: Shell thick, heavy; triangular to rhomboidal, slightly to moderately inflated, often slightly elongated or “humped” in profile; beaks large, flattened, sculpture of three to four loops. Posterior ridge indistinct; shell smooth. Pseudocardinal and lateral teeth distinct, large, compressed beak cavity shallow. Periostracum nearly black, rays lacking. Nacre usually purple, rarely salmon or white.

Habitat: Medium-to-large rivers in mud, sand, or gravel.

Similar species: mucket, spike




6.1" Length: Up to 6"

Spike

Eurynia dilatata (Rafinesque, 1820)

Status: Common

Shell characteristics: Shell moderately thick and solid; elongate-ellipsoid, and slightly laterally compressed; beaks depressed, flattened, sculpture of two to three coarse ridges sometimes drawn up in the middle. Posterior ridge indistinct, rounded; shell smooth. Pseudocardinal teeth well developed and heavy; lateral teeth well developed, short, and straight; beak cavities shallow. Periostracum dark and smooth, rays usually absent. Nacre white, salmon, or purple.



Habitat: Small creeks to large rivers in mud, gravel, and in lakes.

Similar species: Ouachita kidneyshell, black sandshell, elephantear



3.2"


Length: Up to 6"

Wabash pigtoe

Fusconaia flava (Rafinesque, 1820)

Status: Common

Shell characteristics: Shell thin to moderately solid; broadly to sharply triangular, compressed in headwaters with reduced sulcus, thick and inflated with deep sulcus in large river form; beaks depressed, turned forward, sculpture of two to three small ridges. Posterior ridge indistinct; sulcus present. Hinge teeth well developed, with short laterals; deep beak cavity. Periostracum tan to black, rays seldom obvious. Nacre white, sometimes pink to orange, foot usually orange, sometimes white.



Habitat: Creeks to large rivers in gravel and sand. Specimens from the Neosho River basin are of uncertain identification and may represent a separate species.

Similar species: hickorynut, round pigtoe, Ohio pigtoe, deertoe



2.0"


Length: Up to 5"

Ozark pigtoe

Fusconaia ozarkensis (Call, 1887)

Status: Common, but restricted

Shell characteristics: Shell uniformly thin to moderately thick, strong; quadrate to slightly elongate or triangular; beaks low, turned forward, sculpture obscure. Posterior ridge indistinct, with no sulcus; often a pair of diverging grooves on the dorsal slope. Pseudocardinal and lateral teeth well developed and solid; beak cavity moderately deep. Periostracum has a satiny or clothlike sheen; fine green rays may be present. Nacre white to pale pink.



Habitat: Small streams in gravel substrate or mixture of sand and gravel. Only occurs in south flowing Ozark streams of the White and Black river systems.

Similar species: Wabash pigtoe, round pigtoe



2.1"

Length: Up to 4"

Sheepnose


Plethobasus cyphus (Rafinesque, 1820)

Status: Federal Endangered
State Endangered
Species of Conservation Concern

Shell characteristics: Shell moderately thick, heavy, solid; elongate to oval; beak sculpture of two heavy ridges. Posterior ridge weak to moderately developed; shell smooth except for a single row of low, undulating knobs radiating from the umbo to the posterior ventral margin. Pseudocardinal and lateral teeth well developed and heavy, beak cavity shallow. Periostracum yellowish to brownish, rayless, and polished; foot typically bright orange. Nacre white.

Habitat: Medium-to-large rivers in gravel or gravel mixed with sand.

Similar species: mucket, threehorn wartyback, wartyback



3.4" **Length:** Up to 5"

Round pigtoe


Pleurobema sintoxia (Rafinesque, 1820)

Status: Common

Shell characteristics: Shell thick, solid; nearly triangular to nearly round, moderately inflated or compressed; beaks slightly elevated and directed forward, sculpture of two to three small ridges. Posterior ridge indistinct; sulcus absent. Pseudocardinal teeth heavy, lateral teeth well-developed; beak cavity moderately deep. Periostracum dark, generally without defined rays. Nacre white, pink, or orange.

Habitat: Medium-to-large rivers in gravel or gravel-mud substrate.

Similar species: Wabash pigtoe, hickorynut, Ozark pigtoe, Ohio pigtoe, ebonyshell



1.9"


Length: Up to 5"

Ebonyshell

Reginaia ebenus (Lea, 1831)

Status: State Endangered
Species of Conservation Concern

Shell characteristics: Shell thick, solid; subrotund, inflated; beaks inflated, incurved, and projecting forward, sculpture obscure. Posterior ridge indistinct; sulcus absent. Lateral and pseudocardinal teeth well developed and parallel; beak cavity very deep. Periostracum rayless, light brown in young shells, becoming dark brown to black in older individuals.



Habitat: Medium-to-large rivers in fine to coarse gravel.

Similar species: hickorynut, Wabash pigtoe, round pigtoe, Ohio pigtoe




3.1" Length: Up to 6"

Pondhorn

Unio merus tetralasmus (Say, 1831)

Status: Common

Shell characteristics: Shell thin to moderately solid; elongate, rounded anteriorly and pointed posteriorly; moderately inflated; smooth; beaks elevated, sculpture of five to six coarse, concentric semicircles that open anteriorly; usually has two slightly divergent but parallel lines or creases along the posterior ridge. Posterior ridge long and widely rounded. Pseudocardinal teeth distinct and thin, lateral teeth delicate; beak cavities shallow. Periostracum greenish brown and rayless. Nacre white, occasionally salmon.



Habitat: Lakes, ponds, small streams in mud; prefers still water areas.

Similar species: yellow sandshell, spike, giant floater, creeper, cylindrical papershell, tapered pondhorn



3.2"

Length: Up to 5"

Purple wartyback

Cyclonaias tuberculata


(Rafinesque, 1820)

Status: Common

Shell characteristics: Shell thick, heavy, solid; round to quadrate, relatively compressed laterally, with small wing on posterior ridge; beak sculpture of several fine wavy lines. Posterior ridge indistinct; posterior half of shell sculptured with numerous pustules and short ribs. Massive pseudocardinal and lateral teeth and a deep, compressed beak cavity. Periostracum brown to black, lacking rays. Nacre usually purple to deep purple, occasionally coppery, sometimes white.

Habitat: Medium-to-large rivers, primarily in gravel substrates; often found in swift water.

Similar species: pimpleback



3.3"

Length: Up to 4"

Washboard


Megaloniaias nervosa
(Rafinesque, 1820)

Status: Common

Shell characteristics: Shell thick, heavy, solid; elongate-quadrate, compressed to slightly inflated; large; beak sculpture of well-developed double loops. Posterior ridge indistinct; posterior slope fluted; umbonal region covered with zigzag ridges; ridges, vertical creases, wrinkles, and sculpturing present anterior to umbo, and may be present at the ventral margin. Moderately heavy lateral teeth and rather small pseudocardinal teeth for its size. Periostracum brown to black and rayless. Nacre pearly white.

Habitat: Primarily large rivers with gravel or gravel-mud substrates, often in swift water.

Similar species: threeridge, rock pocketbook, bankclimber, pistolgrip, rockshell




— 4.6" — **Length:** Up to 8"

Wartyback

Pustulosa nodulata (Rafinesque, 1820)

Status: Species of Conservation Concern

Shell characteristics: Shell thick, heavy; rounded to quadrate, moderately inflated; beaks prominent, sculpture obscure. Posterior ridge indistinct; two weakly defined rows of knobs diverging from the umbo to the ventral margin of the shell; knobs often are distinctly darker than the rest of the shell; no sulcus. Pseudocardinal and lateral teeth distinct. Periostracum straw colored in young, turning tan to brown with age, rays are lacking. Nacre pearly white.



Habitat: Medium-to-large rivers in sand to fine gravel.

Similar species: pimpleback, mapleleaf, purple wartyback



1.8" Length: Up to 3"

Pimpleback

Pustulosa pustulosa (Lea, 1831)


Status: Common

Shell characteristics: Shell thick, heavy; round to triangular, compressed; beak sculpture obscure. Posterior ridge indistinct; numerous randomly distributed pustules variously developed (may be absent in some individuals or populations), anterior part of shell smooth.

Pseudocardinal and lateral teeth distinct; beak cavity deep. Periostracum brown, often with prominent green ray from umbo halfway down shell. Nacre pearly white.

Habitat: Small streams to large rivers, found in a variety of substrates.

Similar species: wartyback, purple wartyback, rockshell




1.9" Length: Up to 4"

Winged mapleleaf

Quadrula fragosa (Conrad, 1835)

Status: Federal Endangered
State Endangered
Species of Conservation Concern



Shell characteristics: Shell moderately thick; quadrate, with a pronounced wing posterior to the umbo; beak sculpture similar to adult shell. Two rows of pustules from umbo to ventral margin, separated by sulcus. Wing has radiating rows of tubercles, knobs, or flutings. Pseudocardinal and lateral teeth well developed; lateral teeth straight, nearly parallel to ventral margin. Periostracum light tan to dark brown, occasionally with faint green rays. Nacre pearly white to iridescent.

Habitat: Medium-to-large rivers, in gravel or gravel-sand mixtures.

Similar species: mapleleaf, Gulf mapleleaf, wartyback



2.9"

Length: Up to 4"

Gulf mapleleaf


Quadrula nobilis (Conrad, 1854)

Status: Undetermined. Known from the Osage River basin and historically the Grand River, possibly occurs in other areas.

Shell characteristics: Shell thick, heavy; quadrate to ovate to elongate; beaks only slightly above the hinge, sculpture similar to adult. Broad, shallow sulcus present; with two rows of horizontally flattened, elongate pustules on either side of the sulcus, which is often free of pimples or pustules; pustules also occur in areas anterior and posterior to the sulcus. Pseudocardinal teeth erect, heavy, triangular, lateral teeth distinct, straight to slightly curved. Periostracum brown to black. Nacre white.

Habitat: Large rivers in gravel substrates.

Similar species: mapleleaf, winged mapleleaf, pimpleback, pistolgrip



2.7" Length: Up to 5"

Mapleleaf

Quadrula quadrula (Rafinesque, 1820)


Status: Common

Shell characteristics: Shell moderately thick to heavy; quadrate; compressed to inflated (depends on habitat—more inflated in softer substrates); beak sculpture similar to adult shell. Two divergent rows of pustules from umbo to ventral margin, strong sulcus present between rows.

Pseudocardinal and lateral teeth distinct and massive. Periostracum greenish to brown, often with green rays (may be lacking). Nacre pearly white.

Habitat: Medium-to-large rivers, with gravel or sand substrates intermixed with mud; can also be found in reservoirs.

Similar species: winged mapleleaf, Gulf mapleleaf, pimpleback, wartyback, rockshell



2.9"


Length: Up to 5"

Pistolgrip

Quadrula verrucosa (Rafinesque, 1820)

Status: Common

Shell characteristics: Shell heavy, solid; elongate to quadrate or trapezoidal, posterior half tapered; beak sculpture similar to adult shell. Posterior ridge distinct, with large knobs usually on posterior ridge, flutings and pustules prominent especially anterior to ridge; sexually dimorphic — female posterior expanded and compressed. Pseudocardinal teeth large, lateral teeth elongate; shallow beak cavity. Periostracum green to dark brown. Nacre white.



Habitat: Small streams to large rivers, found in a variety of substrates.

Similar species: Gulf mapleleaf, rabbitsfoot, rockshell




3.5"

Length: Up to 8"

Rabbitsfoot

Theliderma cylindrica (Say, 1817)

Status: Federal Threatened
State Endangered
Species of Conservation Concern



Shell characteristics: Shell thick, heavy; elongate to nearly cylindrical, compressed to moderately inflated; beaks only slightly raised above the hinge line, sculpture of coarse folds extending as small tubercles.

Prominent posterior ridge with large nodules or knobs, often sculpturing is below and anterior to the umbos; there is a distinct indentation in the posterior shell margin between the siphons. Well-developed teeth, moderately deep beak cavity. Periostracum typically marked with dense, distinctive greenish chevrons. Nacre white.

Habitat: Medium-to-large rivers in gravel.

Similar species: rockshell, pistolgrip



2.6" Length: Up to 5"

Rockshell


Theliderma metanevra
(Rafinesque, 1820)

Status: Common

Shell characteristics: Shell thick, heavy; quadrate, moderately inflated, with distinct indentation in the posterior shell margin; beak sculpture similar to adult. Very prominent, often slightly concave, posterior ridge with one row of large tubercles extending from umbo to the posterior ventral margin, the remaining shell sculptured with ribs or pustules. Pseudocardinal teeth heavy, lateral teeth short, deep beak cavity. Distinctive inverted greenish chevrons usually present. Nacre pearly white.

Habitat: Prefers gravel substrates in medium-to-large rivers.

Similar species: mapleleaf, pimpleback, Gulf mapleleaf, rabbitsfoot



2.6" Length: Up to 5"



ADDITIONAL SPECIES

POSSIBLY OCCURRING
IN MISSOURI

Ohio pigtoe

Pleurobema cordatum (Rafinesque, 1820)

Status: Unknown. Reported from the St. Francis River.

Shell characteristics: Shell thick, solid; triangular, moderately inflated; beaks elevated above the hinge line and projected forward, sculpture of elevated concentric ridges. Posterior ridge indistinct; shell smooth, with a broad, shallow but distinct sulcus. Pseudocardinal and lateral teeth heavy, lateral teeth straight or nearly so; beak cavity deep. Periostracum dark, without rays. Nacre white, rarely pink or rose-colored.

Habitat: Large rivers in gravel-sand substrates.

Similar species: round pigtoe, ebonyshell, Wabash pigtoe



3.0"

Length: Up to 5"

Tapered pondhorn

Uniomerus declivis (Say, 1831)

Status: Unknown. Reported from the lower St. Francis River.

Shell characteristics: Shell thin, moderately solid; elongate to subrhomboidal, rounded anteriorly and strongly pointed posteriorly; moderately inflated; smooth; beaks slightly elevated, sculpture of five to six coarse, concentric semicircles that open anteriorly. Posterior ridge high and rounded. Pseudocardinal teeth distinct and thin, lateral teeth delicate; beak cavity shallow. Periostracum dark tan to brown. Nacre bluish white.

Habitat: Ponds, lakes, small creeks in fine sand and mud; prefers still water areas.

Similar species: yellow sandshell, spike, giant floater, creeper, cylindrical papershell, pondhorn



— 3.8" — **Length:** Up to 5"


INVASIVE SPECIES

Chinese basket clam

Corbicula fluminea (Müller, 1774)

Status: Invasive

Shell characteristics: Shell relatively strong; rounded to triangular, moderately inflated; beaks moderately inflated and slightly elevated, sculpture of concentric lines, which continue as raised rings. Posterior ridge indistinct; shell with well defined, raised rings. Pseudocardinal teeth triangular and centrally located, two sets of lateral teeth long and straight; beak cavity deep. Periostacum yellowish or brown, often green, becoming darker with age. Nacre purple, white, bluish-white, or purple changing to white. Other species of *Corbicula* may occur in Missouri.



Habitat: Small-to-large rivers and streams, lakes, ponds, in a variety of substrate types.

Similar species: fingernail clams, pea clams




1.8" Length: Up to 2"

Zebra mussel

Dreissena polymorpha (Pallas, 1771)

Status: Invasive

Shell characteristics: Shell thin; elongate triangular, inflated. Dorsal margin straight to broadly rounded and sharp; posterior ridge sharply angulate; anterior end sharply pointed; growth lines not prominent. Pseudocardinal and lateral teeth absent; beak cavity deep, with a small shelf or septum anteriorly. Periostracum highly variable, but most cream or white colored with characteristic dark brown or black bands. Nacre white, polished. Quagga mussel may also occur in Missouri.



Habitat: Small creeks to large rivers, lakes, reservoirs; attaches to virtually any hard surface with glue-like byssal threads.

Similar species: quagga mussel



0.7" Length: Up to 2"

Glossary

Alate winged; having dorsal wing-like structures

Angulate having either the anterior or posterior margins forming a relatively acute angle

Anterior toward the head or front end of an animal

Beak the raised or inflated portion of a bivalve shell, centrally or anteriorly placed along the dorsal margin of the shell

Beak cavity a cavity located inside the shell that extends into the beak

Beak sculpture the raised loops, ridges, or bumps on the beak

Biangulate having two angles

Chevron a V-shaped marking

Compressed flattened or pressed together laterally

Conglutinates small structures made up of gelatinous material and unfertilized eggs that enclose large numbers of glochidia

Dorsal toward the back or top of an animal

Elliptical having the shape of an ellipse or a flattened oval

Emarginate having the margin notched, as in a leaf

Fluted having grooves, furrows, or corrugations on the shell

Glochidia the larval form of unionacean freshwater mussels before metamorphosis

Growth lines darkened lines on the surface of the shell indicating periods of rest during growth

Hinge layers of periostracum that function to keep the two valves of the shell together

Inflated swollen in appearance

Knob a prominent rounded protuberance

Lateral teeth the elongated, raised, and interlocking structures located dorsally along the hinge of the inside of the valves of freshwater mussels

Mussel beds areas containing a high concentration and diversity of mussels

Nacre the inner, lustrous “mother of pearl” layer of the shell, which is often iridescent or colored

Nodule small knot, protuberance, or knob

Ovate (Oval) having a shape with the broader end being the posterior

Periostracum the thin, outside layer or covering of the shell, aka “epidermis”

Posterior rear or toward the back end of an animal

Posterior ridge, posterior slope a ridge or slope emanating from the beak and extending posteriorly to the margin of the shell

Pseudocardinal teeth the triangular, often serrated, molariform structures located on the upper part of the shell in freshwater mussels

Pustule a pimple-like bump or small raised knob on the outside surface of the shell

Pyramidal of or pertaining to a pyramid; triangular

Quadrangle square or rectangular

Rhomboid a parallelogram with opposite sides equal

Sub somewhat or nearly; prefix usually attached to shell outline, as in “subquadrangle” or “subcircular”

Sulcus a shallow depression or furrow

Trapezoidal like a four-sided figure with two parallel sides

Truncate square or broad at end, as if cut off transversely

Tubercle a small, rounded projection or knob

Umbo (Beak) the raised or domed part of the dorsal margin of the shell

Valve the left or right half of a bivalve shell such as a freshwater mussel

Ventral toward the underside or bottom of an animal

Wing (Ala) a dorsal, wing-like structure; is often eroded or broken off

References

- Buchanan, A.C. 1980. Mussels (naiades) of the Meramec River Basin, Missouri. Aquatic Series 17, Missouri Department of Conservation, Jefferson City.
- Cummings, K.S. and C.A. Mayer. 1992. Field guide to freshwater mussels of the Midwest. Illinois Natural History Survey, Manual 5.
- Howells, R.G., R.W. Neck, and H.D. Murray. 1996. Freshwater mussels of Texas. Texas Parks and Wildlife Department, Austin.
- Oesch, R.D. 1995. Missouri naiades: A guide to the mussels of Missouri. Missouri Department of Conservation, Jefferson City.
- Parmalee, P.W. and A.E. Bogan. 1998. The freshwater mussels of Tennessee. University of Tennessee Press, Knoxville.
- Sietman, B.E. 2003. Field guide to the freshwater mussels of Minnesota. Minnesota Department of Natural Resources, St. Paul.
- Utterback, W.I. 1915–1916. The naiades of Missouri. American Midland Naturalist 4(3):41–53; 4(4):97–152; 4(5):181–204; 4(6):244–273; 4(7):311–327; 4(8):339–354; 4(10):432–464.
- Utterback, W.I. 1917. Naiadogeography of Missouri. American Midland Naturalist 5(1):26–30.
- Watters, G.T. 1995. A guide to the freshwater mussels of Ohio. Revised 3rd edition. Ohio Division of Wildlife, Columbus.

Index

Alasmidonta marginata, 17
Amblema plicata, 28
Anodontoides ferussacianus, 18
Arcidens confragosus, 19
Arkansas brokenray, 39

Bankclimber, 49
Black sandshell, 44
Bleedingtooth mussel, 64
Bleufer, 55
Butterfly, 31

Cambarunio hesperus, 29
Canary kingshell, 40
Chinese basket clam, 88
Corbicula fluminea, 88
Creeper, 24
Cumberlandia monodonta, 16
Curtis pearlymussel, 32
Cyclonaias tuberculata, 73
Cylindrical papershell, 18
Cyprogenia aberti, 30

Deertoe, 62
Dreissena polymorpha, 89

Ebonyshell, 71
Elephantear, 65
Elktoe, 17
Ellipsaria lineolata, 31
Ellipse, 63
Elliptio crassidens, 65
Eurynia dilatata, 66
Epioblasma curtisii, 32
Epioblasma triquetra, 33

Fat pocketbook, 51
Fatmucket, 41
Fawnsfoot, 61

Flat floater, 27
Flutedshell, 20
Fragile papershell, 52
Fusconaia flava, 67
Fusconaia ozarkensis, 68

Giant floater, 22
Gulf mapleleaf, 78
Hickorynut, 47
Higgins eye, 37

Lampsilis abrupta, 34
Lampsilis brittsi, 35
Lampsilis cardium, 36
Lampsilis higginsii, 37
Lampsilis rafinesqueana, 38
Lampsilis reeveiana, 39
Lampsilis sietmani, 40
Lampsilis siliquoidea, 41
Lampsilis teres, 42
Lasmigona costata, 20
Leaunio lienosus, 43
Ligumia recta, 44
Lilliput, 59
Little spectaclecase, 43

Mapleleaf, 79
Megaloniais nervosa, 74
Missouri brokenray, 35
Mucket, 48

Neosho mucket, 38
Northern brokenray, 35

Obliquaria reflexa, 45
Obovaria arkansasensis, 46
Obovaria olivaria, 47
Ohio pigtoe, 84

Ortmanniana ligamentina, 48
Ouachita kidneyshell, 56
Ozark pigtoe, 68

Paper pondshell, 26
Pimpleback, 76
Pink heelsplitter, 50
Pink mucket, 34
Pink papershell, 54
Pistolgrip, 80
Plain pocketbook, 36
Plectomerus dombejanus, 49
Plethobasus cyphus, 69
Pleurobema cordatum, 84
Pleurobema sintoxia, 70
Pondhorn, 72
Pondmussel, 57
Potamilus alatus, 50
Potamilus capax, 51
Potamilus fragilis, 52
Potamilus leptodon, 53
Potamilus ohioensis, 54
Potamilus purpuratus, 55
Pterosyna complanata, 21
Ptychobranhus occidentalis, 56
Purple lilliput, 58
Purple wartyback, 73
Pustulosa nodulata, 75
Pustulosa pustulosa, 76
Pyganodon grandis, 22

Quadrula fragosa, 77
Quadrula nobilis, 78
Quadrula quadrula, 79
Quadrula verrucosa, 80

Rabbitsfoot, 81
Reginaia ebenus, 71
Rock pocketbook, 19
Rockshell, 82
Round pigtoe, 70

Sagittunio subrostratus, 57

Salamander mussel, 23
Scaleshell, 53
Sheepnose, 69
Simpsonaias ambigua, 23
Slippershell mussel, 25
Snuffbox, 33
Southern hickorynut, 46
Spectaclecase, 16
Spike, 66
Strophitus undulatus, 24
Sulcularia viridis, 25

Tapered pondhorn, 85
Texas lilliput, 60
Theliderma cylindrica, 81
Theliderma metanevra, 82
Threehorn wartyback, 45
Threeridge, 28
Toxolasma lividum, 58
Toxolasma parvum, 59
Toxolasma texasiense, 60
Truncilla donaciformis, 61
Truncilla truncata, 62

Uniomerus declivis, 85
Uniomerus tetralasmus, 72
Utterbackia imbecillis, 26
Utterbackiana suborbiculata, 27

Venustaconcha ellipsiformis, 63
Venustaconcha pleasii, 64

Wabash pigtoe, 67
Wartyback, 75
Washboard, 74
Western fanshell, 30
Western rainbow, 29
White heelsplitter, 21
Winged mapleleaf, 77

Yellow sandshell, 42

Zebra mussel, 89



Serving nature and you®

